FOOD ALLERGY / INSECT ALLERGY ACTION PLAN

	◆ STEP 1: TREAT	
	SYMPTOMS:	GIVE CHECKED MEDICATION(S)**: ** (TO BE DETERMINED BY PHYSICIAN AUTHORIZING TREATMENT)
	 If food allergen has been ingested, or child has been stung, but no symptoms: 	Epinephrine Antihistamine
	Mouth: Itching, tingling, or swelling of lips, tongue, mouth	Epinephrine Antihistamine
	Skin: Hives, itchy rash, swelling of the face or extremities	Epinephrine Antihistamine
	Gut: Nausea, abdominal cramps, vomiting, diarrhea	Epinephrine Antihistamine
	Throat †: Tightening of throat, hoarseness, hacking cough	Epinephrine Antihistamine
	Lung † : Shortness of Breath, repetitive coughing, wheezing	Epinephrine Antihistamine
	 Heart †: Weak or thready pulse, low blood pressure, fainting, pale, blueness 	Epinephrine Antihistamine
	Other †:	Epinephrine Antihistamine
	 If reaction is progressing (several of the above areas affected), give: 	Epinephrine Antihistamine
	†Potentially life-threatening. The severity of DOSAGE	symptoms can quickly change.
inephr tihista	rine: iniect intramuscularly (circle one) EpiPen® EpiPer	n® Jr. Twinject® o.3mg Twinject® o.15mg
her: Gi	mine: Givemedication / dose / re	oute
	medication / dose / ro MPORTANT: ASTHMA INHALERS AND / OR ANTIHISTAM EPINEPHRINE IN ANA STEP 2: EMERGEN	oute INES CANNOT BE DEPENDED ON TO REPLAC PHYLAXIS
	Call 911 (or Rescue Squad). State that an allergic re be needed.	action has been treated, and additional epinephri
2. I	Dr Phone Number:	
	ParentPhone Number:	
	Emergency Contacts: (Name / Relationship) 1) 1) D 1)	

(REQUIRED)